


WHAT IS CLAIMED IS:

1. A block polymer comprising:
 - i) one or more cationic group-containing units; and
 - 5 ii) optionally one or more additional building block units;
provided that the block polymer has an average cationic charge density of about 15 or less.
2. The block polymer according to Claim 1 wherein said block polymer further comprises:
 - 10 iii) one or more units having one or more hydroxyl groups,
provided that said polymer has a Hydroxyl Group Density of about 0.5 or less.
3. The block polymer according to Claim 1 wherein said block polymer further 15 comprises:
 - iv) one or more units having one or more hydrophobe groups selected from the group consisting of non-hydroxyl groups, non-cationic groups, non-anionic groups, non-carbonyl groups, and/or non-H-bonding groups.
- 20 4. The block polymer according to Claim 2 wherein said block polymer has a Hydroxyl Group Density of from about 0.0001 to about 0.4.
5. The block polymer according to Claim 1, wherein said polymer further comprises:
 - 25 v) units capable of having an anionic charge at a pH of from about 4 to about 12;
 - vi) units capable of having an anionic charge and a cationic charge at a pH of from about 4 to about 12;
 - vii) units having no charge at a pH of from about 4 to about 12; and
 - 30 viii) mixtures of units (v), (vi), (vii), and (viii).
6. The block polymer according to Claim 5, wherein said polymer average cationic charge density ranges from about 0.05 to about 5 units per 100 daltons molecular weight at a pH of about 4 to about 12.


7. The block polymer according to Claim 1, wherein said polymer has an average molecular weight of from about 1,000 to about 2,000,000 daltons.

8. The block polymer according to Claim 1, wherein said block polymer comprises a cationic unit of the formula:



wherein each of R¹, R² and R³ are independently selected from the group consisting of hydrogen, C₁ to C₆ alkyl, and mixtures thereof; T is selected from the group consisting of substituted or unsubstituted, saturated or unsaturated, linear or branched radicals selected from the group consisting of alkyl, cycloalkyl, aryl, alkaryl, aralkyl, heterocyclic ring, silyl, nitro, halo, cyano, sulfonato, alkoxy, keto, ester, ether, carbonyl, amido, amino, glycidyl, carbanato, carbamate, carboxylic, and carboalkoxy radicals and mixtures thereof; Z is selected from the group consisting of: -(CH₂)-, (CH₂-CH=CH)-, -(CH₂-CHOH)-, (CH₂-CHNR⁴)-, -(CH₂-CHR⁵-O)- and mixtures thereof; R⁴ and R⁵ are selected from the group consisting of hydrogen, C₁ to C₆ alkyl and mixtures thereof; z is an integer selected from about 0 to about 12; A is NR⁶R⁷ or NR⁶R⁷R⁸ wherein each of R⁶, R⁷ and R⁸, when present, are independently selected from the group consisting of H, C₁-C₈ linear or branched alkyl, alkyleneoxy having the formula:


20








wherein R⁹ is C₂-C₄ linear or branched alkylene, and mixtures thereof; R¹⁰ is hydrogen, C₁-C₄ alkyl, and mixtures thereof; and y is from 1 to about 10.

25


9. The block polymer according to Claim 1, wherein said block polymer comprises a cationic unit of the formula selected from the group consisting of:








5



10


10. The block polymer according to Claim 1 wherein the block polymer is selected from the group consisting of:

15




20

25




30

5



10



15




20

25

30



35



wherein R¹ is selected from H and CH₃; R² is selected from H and SO₃H; x represents the total number of monomer units within the block polymer; m, n, o, p, q when present, represent the mole ratio of their respective monomeric units in a given block polymer where at least two different monomeric units are present in the block polymer.

11. A block polymer comprising at least a first homopolymeric unit comprising a series of first cationic monomeric units and at least a second homopolymeric unit comprising a series of second polymeric units,

5 at least said first monomeric units capable of having a cationic charge at a pH of from about 4 to about 12;

provided that said polymer has an average cationic charge density from about 0.05 to about 5 units per 100 daltons molecular weight at a pH of from about 4 to about 12.

10 12. A block polymer comprising at least one homopolymeric block of monomeric units A and at least one member of the group consisting of a homopolymeric block of monomeric units B and a homopolymeric block of monomeric units C:

A. said block of cationic monomeric units A having a Formula I:


15





wherein R^1 is H or an alkyl having 1 to 10 carbon atoms,

20


R^2 is a moiety selected from the group consisting of



5




, and



wherein R^3 is selected from the group consisting of


10



a is an integer from 0 to 16; b is an integer from 2 to 10; c is an integer from 2 to 10; d is an integer from 1 to 100;

15

R^4 and R^5 are independently selected from the group consisting of -H, and





20

R^8 is independently selected from the group consisting of a bond and an alkylene having 1 to 18 carbon atoms;

R^9 and R^{10} are independently selected from the group consisting of -H, alkyl having 1 to 10 carbon atoms;


25

R^{12} and R^{13} are independently selected from the group consisting of H and alkyl having from 1 to 10 carbon atoms;



wherein t is an integer from 2 to 10;


B. said monomeric unit B is selected from the group consisting of:
15 a monomeric unit of Formula IV




20


wherein R^{20} is selected from the group consisting of H and CH_3 ;
 R^{21} is selected from the group consisting of:

25







5




10




wherein e is an integer from 3 to 25;

15



wherein f is an integer from 0 to 25;

20



25


wherein g is an integer from 1 to 100,

h is an integer from 1 to 100,



R²³ is -H, -CH₃ or -C₂H₅,

R²⁴ is -CH₃ or -C₂H₅;


30



wherein j is an integer from 1 to 25;




15 wherein k is an integer from 1 to 25;




-NH-(CH₂)_r-NH₂·HCl, wherein r is an integer from 1 to 25; and

a polyhydroxy monomeric unit of Formula VI:




wherein w is an integer from 1 to 50; and

C. monomeric unit C is selected from the group consisting of:



wherein R²⁵ is -H or -CH₃,

5



10

wherein R²⁶ is -H or CH₃, and

x represents the total number of monomeric units within the block polymer; m, n, o, when 15 present, represent the mole ratio of their respective monomeric units in a given block polymer, wherein at least two different monomeric units are present in the block polymer.

13. The block polymer of Claim 12, wherein m is greater than 1, n is greater than 1 and o is greater than 1.


20

14. The block polymer of Claim 12, wherein said polymer comprises at least one said monomeric unit A, at least one said monomeric unit B and at least one said monomeric unit C.

15. The block polymer of Claim 12, wherein said at least one monomeric unit A is selected from the group consisting of:

5

10



wherein R³⁰ is H or -CH₃,


wherein R³¹ is a bond or $\text{C}=\text{O}$, and

15

R³² and R³³ are -CH₃ or -C₂H₅.

20


25



wherein R³⁸ is selected from the group consisting of H and CH₃ and

R⁴⁰ is selected from the group consisting of -CH₂CH₂-OH and

30




and isomers thereof,

said terpolymer comprising said at least one monomeric unit C,

wherein the molar ratio of said monomeric unit A : monomeric unit B : monomeric unit C is 1 to 9 : 1 to 9 : 1 to 6 respectively.

17. The block polymer of Claim 12, wherein the at least one monomeric unit B has
5 the formula:

10




wherein q ranges from 1 to 12.

18. The block polymer of Claim 17, wherein the polymer is a terpolymer, said at least one monomeric unit A is selected from the group consisting of:

15

20



wherein R¹⁰ is H or CH₃,


R¹¹ is a bond or $\text{---C=---}^{\text{O}}$, and R¹² and R¹³ are -CH₃ or -C₂H₅, and said monomer comprises said at least one monomeric unit C.

5

19. The polymer of Claim 18, wherein the molar ratio of monomeric unit A : monomeric unit B : monomeric unit C ranges from 1 to 9 : 1 to 9 : 1 to 3 respectively.

20. The block polymer of Claim 12, wherein said at least one monomeric unit A has a
10 formula selected from the group consisting of:


15




20

21. The block polymer of Claim 12, wherein said at least one monomeric unit B has a formula selected from the group consisting of:

25




30




and

5




22. The block polymer of Claim 12, wherein said at least one monomeric unit B is selected from the group consisting of:

10




wherein n is an integer from 1 to 50, preferably 2 to 30, more preferably 7 to 27,


15




20




25




30



5



10



15

23. The block polymer of Claim 12, selected from the group consisting of:
 poly(HEA-co-DMAM-co-AA) terpolymer,
 poly(HPA-co-DMAM-co-AA) terpolymer, and
 poly(PEG-acrylate-co-DMAM-co-AA) terpolymer.

20

24. The polymer of Claim 12, selected from the group consisting of:
 poly(HEA-co-DMAM) copolymer, poly(DMAM-co-butylvinylether) copolymer
 and poly(2-diethylaminoethylvinyl ether-co-ethyleneglycol monovinyl ether).

25

25. A method for cleaning hair or skin comprising applying an effective amount of a
 cleaning composition comprising the polymer of Claim 1 and at least one deterotive surfactant to
 hair or skin in need of cleaning, provided that a 10% aqueous solution of said composition has a
 pH from about 4 to about 9.

30

26. The method of Claim 25, wherein said composition further comprises at least one
 member of the group consisting of a pearlizing agent, a silicone hair conditioning agent, and an
 antidandruff ingredient.

27. The method of Claim 26, wherein said composition comprises:

- a) said pearlizing agent
- b) a nonionic surfactant
- c) an amphoteric surfactant
- 5 d) a glycol emulsifier
- e) water.

28. The method of Claim 26, wherein said composition comprises at least one amphoteric surfactant and said amphoteric surfactant comprises at least one member of the group 10 consisting of:

the alkali salts of alkyl amphodipropionates, alkyl amphodiacetates, alkyl amphoglycinates, alkyl amphopropyl sulfonates and alkyl amphopropionates wherein alkyl represents an alkyl group having 6 to 20 carbon atoms.

15 29. The method of Claim 28, wherein in said at least one amphoteric surfactant the alkyl group is derived from coconut oil or is a lauryl group.

30. A method for cleaning hair or skin comprising applying an effective amount of a 20 cleaning composition comprising the polymer of Claim 13 and at least one surfactant to hair or skin in need of cleaning.

31. A composition for cleaning hair or skin comprising:
the block polymer of Claim 1,
at least one detergentsurfactant, and at least one member of the group consisting 25 of a pearlizing agent, a silicone hair conditioning agent, and an antidandruff ingredient, provided that a 10% aqueous solution of said composition has a pH from about 4 to about 12.

32. A composition for cleaning hair or skin comprising:
the block polymer of Claim 12,
30 at least one surfactant, and at least one member of the group consisting of a pearlizing agent, a silicone hair conditioning agent, and an antidandruff ingredient.

33. The composition of Claim 32, wherein said silicone compound is an alpha, omega-trimethylsilyl-polydimethylsioloxane having a viscosity at 25°C of at least 25 centistokes and less than 60,000 centistokes.

5 34. A method for washing a fabric article in a washing medium comprising:
applying an effective amount of a laundry cleaning composition comprising the polymer of Claim 1 and at least one detergent surfactant to a fabric article in need of cleaning.

10 35. The method of Claim 34, wherein said composition washes a colored fabric article.

15 36. The method of Claim 34, wherein said composition comprises at least one member of the group consisting of an aminosilicone, a Gemini surfactant, a detergency builder, a bleach, an activator for percompound bleach, a soil suspending agent, a soil antiredeposition agent, a foam suppressant agent and a fabric softener.

37. The method of Claim 34, wherein said composition comprises a foam suppressant agent.

20 38. A method for washing a fabric article in a washing medium comprising:
applying an effective amount of a laundry cleaning composition the polymer of Claim 13 and at least one detergent surfactant to a fabric article in need of cleaning.

25 39. A composition for washing a fabric article comprising:
the polymer of Claim 1;
at least one detergent surfactant; and
at least one member of the group consisting of an aminosilicone, a Gemini surfactant, a detergency builder, a bleach, an activator for percompound bleach, a soil suspending agent, a soil antiredeposition agent, a foam suppressant agent and a fabric softener.

30 40. A method for extinguishing fire comprising applying a foam to a fire, wherein the foam comprises a foaming agent and a polymer of Claim 1.

41. A method for treating at least one agricultural substrate selected from the group consisting of plants, seeds and soil comprising,

5 applying to the substrate a foam comprising at least one agricultural chemical selected from the group consisting of a herbicide, a pesticide, and a fungicide, a foaming agent and a polymer of Claim 1.

42. A method comprising, injecting into a subterranean formation, a foam comprising a foaming agent and a polymer of Claim 1.

10

43. A method comprising shaving hair from skin comprising applying foam shaving cream to the skin, said shaving cream comprising a foaming agent and a polymer of Claim 1.

15

44. A method comprising shaving hair from skin comprising applying a shaving gel to the skin, said gel comprising a foaming agent and a polymer of Claim 1.

45. A method for removing hair from skin comprising applying a depiliatory foam to skin, said foam comprising a foaming agent and a block polymer of Claim 1.

20

46. A method of cleaning hard bathroom surfaces comprising applying to said surfaces a foam cleaner comprising a foaming agent and a polymer of Claim 1.

25

47. A process for making paper comprising aiding retention of titanium dioxide on the paper during the paper making comprising treating the paper with an aqueous solution comprising titanium dioxide and a polymer of Claim 1.